

Appointment Date:

Appointment Time:

If you have any problems completing the instructions, call: (760) 924-4014

*if calling after clinic hours call (760) 934-3311 and ask for the PACU.

Please follow all instructions carefully

On the day of your colonoscopy, your colon must be clean on the inside, and you must have a ride home following the procedure. An unclean colon can result in your test being rescheduled, missed lesions, increased procedure time, and a potential increase in complication rates. Bring a list of all of your current medications with you and let your doctor know if you have diabetes, kidney problems, or might be pregnant.

7 DAYS BEFORE YOUR COLONOSCOPY

If you are diabetic, you will get special instructions on how to adjust your medication. If you are taking any medication that affects blood clotting, you may have to adjust 7 to 3 days before the test. Be sure to check what medications you should take with your doctor. Examples of prescription blood thinners include Agrylin™, Coumadin® (warfarin), Plavix™ (clopidogrel), Ticlid™ (ticlopidine), Xarelto™ (rivaroxaban), Pradaxa® (dabigatran), Effient™ (prasugrel), Lovenox™ (enoxaparin), and over-the-counter medications including aspirin and NSAIDs (Advil™/ibuprofen, Motrin™, Aleve®, and Ecotrin™).

3 DAYS BEFORE YOUR COLONOSCOPY

- Stop eating any nuts, seeds, corn, or popcorn.

2 DAYS BEFORE YOUR COLONOSCOPY

- Drink 8 glasses of water or other liquids during the day to avoid dehydration.



THE DAY BEFORE YOUR COLONOSCOPY

- **STOP eating all solid foods**
- **START your clear liquid diet.** (No red, blue, or purple flavors.) Clear liquids mean that you can see through them. Gatorade, Pedialyte, or Powerade
- Clear broth or bouillon
- Coffee or tea (no milk or creamer)
- Carbonated and non-carbonated soft drinks, sparkling water, Sprite, 7-Up, strained fruit juices (no pulp) like apple or white grape
- Jell-O, popsicles, and hard candy
- **START THE PREP** that your doctor has checked on this page and follow instructions closely. Your doctor may add other oral laxatives to be used with your prep
- Brand names include Dulcolax, Citroma, Citrate of Magnesia, and LiquaPrep

THE DAY OF YOUR COLONOSCOPY

- Unless your doctor tells you otherwise, you will usually take your morning medication with a sip of water up to 2 hours before your colonoscopy. Diabetic patients will need to adjust their medication as directed.
- You will receive sedation and will need someone to drive you home.



Health Information

For Patients and the Community

Directions for Taking MoviPrep

Dose 1: 4 pm the night before the colonoscopy:

Drink one 8-ounce glass of the solution every 15 to 20 minutes while continuing to drink clear liquids.

FINISH ALL OF THE PREP!

DURING THE PREP: IT IS IMPORTANT THAT YOU REPLACE FLUIDS AND PREVENT DEHYDRATION BY DRINKING LARGE AMOUNTS OF CLEAR LIQUIDS UNTIL TWO HOURS BEFORE THE COLONOSCOPY.

Helpful Tips

1. Stay near a toilet; you will have diarrhea. Ideally, your bowel movements will become either yellow or clear liquid after a complete prep.
2. Side effects of some preps include nausea, abdominal pain, bloating, vomiting, thirst, dizziness, and dehydration. If you feel nauseous or vomit, take a 30 minute break, rinse your mouth, and then continue drinking your prep. If you throw up your bowel prep, just try your best to drink as much bowel prep as you can tolerate and report for your endoscopy as scheduled. Be sure to alert your doctor that you could not tolerate your entire bowel prep.
3. Feeling some abdominal cramping is normal.
4. Petroleum jelly (Vaseline™) around the anus or using baby wipes may decrease skin irritation.

You will get a call the night before your Colonoscopy between 4:00-7:00 to tell you what time to arrive for your procedure.